



A COMPENDIUM ON THE FUNGAL DISEASES OF VEGETABLES IN KOTA DISTRICT OF RAJASTHAN

Dr. Rashmi Pant¹, Lalita Sharma², Ravi P. Shringi³

¹ Department of Basic & Applied Life Science , MIMT, Dadabari Ext. , Kota (Raj.),(India)

^{2,3}Department of Botan , Govt PG College, Kota , Rajasthan,(India)

ABSTRACT

Fungi constitute a major problem in the production, storage and processing of agricultural products especially Vegetable crops. Vegetables belonging to the families cucurbitaceae, brassicaceae and solanaceae are important due to their nutritional as well as economical values. The farmers of Hadoti region face heavy yield losses both in quality and quantity of these crops due to damage as a result of various diseases caused by fungal pathogens. There are different pathogens which were found to be associated with the various angiospermic diseases of vegetables in Kota District . These diseases develop through soil-borne, infections and by insect as a vectors. However, no comprehensive review is available on the fungal disease of vegetables in Kota District . The present review gives inclusive information regarding various pathological aspects of the fungal disease, their causes and management strategies accomplished on them.

Key Words: Nutritional, Angiospermic Diseases, Pathological Aspect, Soil Borne.

I. INTRODUCTION

The population of the world is increasing at an alarming rate. It has been projected to grow to 8.2 billion by the year

2025 and is likely to approach 9.3 billion in 2050 (DESA,

2000). Feeding such rapidly projected growing population is becoming a big problem. India has advantage in the production of a variety of fresh crops especially vegetables during dry season and in wet season also in the low lands (Bessey, E.A. 1950)Estimates of production losses in developing countries are hard to evaluate. Post harvest losses of fruit and vegetables in some Asian countries have been estimated to reach 50% (Tsoho, 2004; FAO,

2008; FAO, 2012) hence minimizing post harvest losses of already produced food is more sustainable than increasing production (Kader and Rolle, 2004). Post-harvest diseases of fruits and vegetables caused by fungal and bacterial pathogens result in significant economic losses. Diseases caused by Microbial pathogens have been a major challenge to agriculture, health as well as national



economy (Shehu et al., 2014^a). In Rajasthan, vegetables are commonly displayed on benches, sacks and in baskets for prospective buyers. While stored in the open markets the produce are susceptible to microbial invasion and colonization leading to contamination (Muhammad *et al.*, 2004; Shehu et al., 2014^b). Vegetable contamination with fungal pathogens leading to diseases development on the produce can also occur due to pre-harvest practices applied on the diseased vegetable crops such as plant fertilization with manure, sewage sludge and from irrigated water. Other aspects which contribute to disease of vegetables include injuries, high temperature and storage in previously contaminated bins (Shehu *et al.*, 2014^c). The study area is Kota district which lies between latitude 13° 3' 49⁰N, longitude 5° 14' 89⁰E and at an altitude of 272 m above the sea level. It is located in the extreme North Western part of India. The soil is predominantly ferruginous tropical type, texturally sandy and pH of the soil ranges between 6 and 7. Rainfall starts late from June and ends early, in September. The highest temperatures of 45°C during the hot season are experienced in the months of April to June. Production of vegetable is a major occupation of people of Rajasthan and neighboring states in north western Nigeria, especially in the dry season when rainy season cultivation is over. This activity is both an income earner as well as a source of employment to a large proportion or otherwise underemployed labour force (Ibrahim, 2005; Salau, 2012; Shehu and Aliero, 2011)..

In current paper a review on the fungal disease of vegetables is described with reference to production, occurrence, symptomology, incidence, pathogenicity, losses and disease management.

II. VEGETABLE PRODUCTION

Our dependence on healthy vegetable crops as reliable source of food traced all barriers of nation and culture. Consumers in Nigeria now demand excellent quality vegetable from the farmers and vendors that produce or sale large volumes of high quality vegetables (Shehu *et al.*,

2014^d). Amaranth (*Amaranthus caudatus*), roselle (*Hibiscus sabdariffa*), kenaf (*Hibiscus cannabinus*) Onion (*Allium cepa*) and tomato (*Lycopersicon esculentum*) are among the most common vegetables cultivated by farmers in Kota district of Rajasthan State. The vegetables grow in varied types of soil – sandy loam to clay and also tolerate moderate acidic and saline soils. The optimum temperature for its cultivation vegetables is 15 to 27⁰C. Abelmoscus esculentus is a quick – growing green vegetable and is often uprooted when it is 8 to 10 cm tall (3 to 4 weeks after sowing). Its flowers mature in 90 to 95 days. It is of the malvaceae family, it bears leaves, follicle, seeds which are edible and have versatile uses. The plant has an overall growing period of 4 to 6 months. The young plant is often uprooted when it is 3 to 4 weeks old for sale and consumption (Salau, 2012).



2.1 Diseases of Vegetables

Fungal disease constitutes a menace in vegetable production, and many pathogens have been associated with vegetable plant in the field as well in the storage and processing stage.

The principal fungal fruit rots reported all over the world with varying intensities on vegetables includes *Alternaria* rot caused by *A. solani* and *A. tenuis*, *Phytophthora* rot caused by *Phytophthora infestans*, *Phytophthora nicotianae* var. *parasitica*, Anthracnose ripe rot caused by *Colletotrichum phomoides*, *Phoma* rot *Phoma destructiva* and *Fusarium* rot caused by *Fusarium* spp. (Ibrahim, 2002; Iqbal *et al.*, 2003; Patel *et al.*, 2005; Ali *et al.*, 2005; Salau *et al.*, 2012). Early blight, caused by the fungus *Alternaria solani*, appears first on the lower leaves, usually after a heavy fruit set. It normally forms a few large (up to 1/2 inch) spots on a leaf. The spots are dark brown to black and roughly circular. Concentric rings develop in the spot forming a "target spot". The leaf area around each target spot turns yellow, and soon the entire leaf turns yellow and drops. *Alternaria* rot has been considered as the most common disease of tomato fruits and causes heavy losses in quality of the fruits, thus rendering large quantity of tomato fruits unfit for consumption. The disease was reported by (MC Kellar M.E. and E.B. Nelson 2003),

The causal organism was isolated from the dead leaves of leafy vegetables and afterward it was reported from the decaying tomato fruits. Salau *et al.*,(2012) reported that *A. solani* was mainly responsible for the tomato decay and typical symptoms occurred on stem were brown to black, depressed and usually with distinct rings. Ibrahim (2002) reported that the spot range in size from minute pin-heads to areas extending completely across the surface of the fruit giving it a flattened appearance. *Fusarium* rot is considered most destructive on vegetables has been reported from (Ibrahim *et al.*,2011). Number of different *Fusarium* species has been found associated with this rot. Fungal species of *Acremonium*, *Alternaria*, *Fusarium*, and *Plectosporium* were also detected in surveyed sites (Salau *et al.*, 2012). Several vegetable crops such as *Solanum nigrum*, *Lycopersicon* , *Cucurbits* are effected by these fungal pathogens .About 62.55% of vegetable crops get affected during the favourable environmental conditions in which they are grown. Hence, this is the biggest reason for total economical loss of the income of farmers belonging Kota region of Rajathan state.

III. SOURCES OF INOCULUMS

Microbial organisms depend on different factors including soil condition that could be the reservoir of food borne pathogens or water for irrigation (Amir H & Alabouvette C 1993). Water may be source of different pathogens; they could be transferred from water to the vegetables Veeken A.H.1987) organic fertilizers (sewage sludge, animal manure, compost) may also be potential risk for contamination of vegetables. Vegetables may also be contaminated during harvest (due to lack of sanitary facilities to workers or dirty storage facility) or post harvest treatment (handling, storage and transportation). Therefore, to minimize the risk of infection or intoxication associated with vegetables, potential source of contamination should be identified and specific



measures and intervention to prevent and minimize the risk of contamination should be considered and correctly implemented (Aliero and Ibrahim 2014).

IV. FACTORS THAT INFLUENCE VEGETABLE DISEASE

Vegetable losses vary each year, prevailing weather while the crops are growing and at harvest contribute greatly to the possibility of decay certain cultivars are more prone to decay than are others to specific pathogens. In a recent study, it was found that resistance of major vegetable cultivars to the fungi that cause blue mold, gray mold, bull's-eye rot, and Mucor rot was dependent on cultivar. Condition of the crop, as determined by fertilizer and soil factors, are very important in susceptibility of the crop to disease. Maturity of the crop at harvest, handling and type of storage has a great deal of influence on how long the crop can be stored without decay. Weather affects many factors related to plant diseases, from the amount of inoculum that overwinters successfully to the amount of pesticide residue that remains on the crop at harvest (Shehu and Muhammad, 2011).

Packing Sanitation it is important to maintain sanitary conditions in all areas where produce is packed. Organic matter (culls, extraneous plant parts, soil) can act as substrates for decay-causing pathogens. For example, in vegetables and pear packinghouses, the flumes and dump tank accumulate spores and may act as sources of Contamination if steps are not taken to destroy or remove them.

V. PATHOGENICITY

Literature revealed that several methods of inoculation have been adopted by different workers for confirming the Pathogenicity of the causal fungi isolated from collected vegetables in Kota region.

According to Shehu and Muhammad (2011) who studied Fungi associated with storage rots of onion bulbs in Sokoto revealed that *Aspergillus niger*, *A. flavus*, *A. fumigatus*, *Alternaria porri*, *Fusarium ox-ysporum*, *Rhizopus stolonifer* and *Penicillium citrinum* were found to be pathogenic to the onion bulbs, these fungi leading to rapid disintegration of the infected bulbs within 21 days of inoculation. The finding from this investigation indicates that the aforementioned fungi were associated with deterioration of onion bulbs. This agrees with the reports of other researchers (Muhammad *et al.*, 2004; Dimka and Onuegbu, 2010) that fungi constitute a menace in the storage of many agricultural commodities including fruits, vegetables and nuts.

Similarly Salau *et al.* (2012) implicate that *Aspergillus* spp., *Alternaria* spp, *Penicillium* spp., *Rhizophus* spp. and *Saccharomyces* spp from there finding on Studies of fungi associated with vegetable plant diseases in Kota region. Metropolis revealed that within a week (7 days), a soft rotten area developed around the inoculation point of tomato, the same species were isolated from the rotten part, while all smearing inoculated leaves of vegetables remained green and symptomatic in all smearing, and un- smeared controls also remained green and asymptomatic in both places. However symptoms were the same as those previously observed in the field in which species of the *Aspergillus* group were identified as the causal agent of disease in these (Spinach, Lettuce, tomato and Cabbage) vegetable



crops. The result obtained is in consonant with the finding of Kutama *et al.*, (2007) who reported several species of Fungal Pathogens found associated with Tomatoes and his storage vessel (*Aspergillus* spp., *Alternaria* spp, *Penicillium* spp., *Rhizopus* spp. and *Saccharomyces* spp). This could be associated with hygienic conditions of the cultivation site of tomatoes.

VI. ECONOMIC IMPLICATION

Losses from post harvest deterioration of vegetables have been substantial during storage, wholesale, retail, and consumer levels. The total losses are very difficult to establish because research has generally considered only one or two levels, and little work has been done to determine losses at different levels of storage, wholesale, retail and consumer level. Nevertheless in Rajasthan state, losses of vegetables are estimated to range from 5% to 43% of the harvested vegetables (Shehu *et al.*,2014^b), most of the vegetable decay results from microbial infection (through wounds made during harvest) and postharvest handling. However, infection may takes place in the orchard during the growing season and remain latent. And in storage, the pathogens become active again and invade the vegetable tissues.

VII. MANAGEMENT STRATEGIES

A variety of techniques have been used to reduce the decay and other forms of vegetable losses in the area, these include heaping of onion on the floor of straw hurt, sanitation to reduce pathogens inoculums from all types of vegetables, careful handling to reduce wounding, storing vegetables at low temperature, treatment of vegetables with fungal inhibitor among others.

VIII. CONCLUSION

It was concluded from the above review that fungal disease of vegetables is caused by several pathogenic fungi under different conditions. Many studies have been carried out with respect to occurrence, causal organisms, severity, losses, Pathogenicity and disease management. This review may help the future researchers to devise a concrete strategy for evaluating different pathological aspects and management of the post harvest fungal diseases of vegetables. However, further study is needed to reveal all the other recent reports about various pathological aspects on the fungal diseases of vegetables and management strategies opted for diseases.

REFERENCES

- [1] Amir H & Alabouvette C 1993. *Involvement of soil abiotic factors in the mechanisms of soil suppressiveness to Fusarium Wilt. Soil Biol. Biochem* 25:157-164.
- [2] Anonymous. 1959. *Proc. Seminar on Diseases of Horticultural Plants* I.C.A.R. New Delhi.
- [3] Ashkar, S.A. and Ries, S.K. 1971. *Journal of the American Society for Horticultural Science* 96:448-452
- [4] Abdullahi K, Abubakar U, Adamu T, Daneji AI, Aliyu RU, Jiya N, Ibraheem MTO, Nata'ala SU (2009). Malaria in Sokoto, North Western Nigeria *Afri. J. Biotechnol.*, **8** (24): 7101-7105,



- [5] Ali S, Rivera VV, Secor GA (2005). First report of *Fusarium graminearum* causing dry rot of potato in North Dakota. *Plant Dis.*, **89**: 105-109.
- [6] Aliero AA, Ibrahim AD (2014). Aflatogenic and Aflatoxin contents of dried vegetables sold in Sokoto Metropolis, Nigeria. *Journal of Emerging Trends in Engineering and Applied Sciences (JETEAS)* 5(7): 116-120
- DESA, Department of Economic and Social Affairs (2000). World Population Prospects: The 2000 revision. United Nation population division, Department of Economic And Social Affairs in Badaurakis. *Int.J. Agribus.*, 18(4): 543-558.
- [7] B.R. 1983 Research needs to upgrade the nutritional quality of common bean (*Phaseles vulgaris*). *Plant Foods Human Nutrition* 32, 101-110.
- [8] Baker, K.F. & R.J. Cook 1974. *Biological Control of Plant Pathogens* pp 433.
- [9] Bessey, E.A. 1950. *Morphology and Taxonomy of Fungi*. The Blakiston Co. , Philadelphia.
- [10] Bruehl GW (Ed) *Biology and Control of Soil Borne Pathogens* (pp 137-144). American Phytopathological Society.
- [11] Chester, K.S. 1950. *Nature and Prevention of Plant Diseases*. 525 pp. Blackiston, Philadelphia, Pennsylvania.
- [12] Chupp, C. and A.F. Sherf. 1960. *Vegetable Diseases and Their Control*. 693 pp. Ronald Press, New York.
- [13] Dimka SON, Onuegbu BA (2010). Mycoflora of copra and effect of brining on some properties of copra in Nigeria. *Agriculture and Biology Journal of North America*: 2151 – 7525
- [14] Dogondaji SD, Baba KM, Muhammad I, Magaji MD (2005). Evaluation of onion storage losses and implication for food security in Sokoto Metropolis. *Bulletin of Science Association of Nigeria*. 26: 10 – 14
- FAO (2008). Food and Agriculture Organization. The World Vegetable Centre Newsletter. www.avrdc.org
- [15] FAO (2012). Food and Agriculture Organization. The State of Food and Agriculture www.fao.org/docrep/016/i3027e/i3027e
- [16] Ibeyessie J (2007). Bacterial pathogens recovered from vegetables irrigated by waste water. *Journal of Environmental health*, **37** (6): 711-718
- [17] Ibrahim AD, Hussaini H, Sani A, Aliero AA, Yakubu SE (2011). Volatile metabolite profiling to discriminate diseases of tomato fruit inoculated with three toxigenic fungal pathogens. *Research in biotechnology*, 2 (3): 14-22
- [18] Iqbal SM, Ghafoor ZA, Haqqani (2003). Pathogenicity and fungicidal efficacy for sclerotinia rot of brinjal. *Int. J. Agric. Biol.*, **5**(4): 618-620.
- [19] Klein E.J. Katan and A. Gamliel 2011. Soil suppressiveness to *Fusarium* disease following *Organic amendment and solarization* *plant disease* 95: 1116-1123
- [20] Lilly, V.G. and H.L. Barnett. 1951. *Physiology of fungi*. McGraw-Hill, New York, studied the habit and habitat of different fungi.
- [21] Lin S & Baker R 1980. *Mechanism of Biological control in soil suppressive to Rhizoctonia solani*. *Phytopathology* 70: 404-412.



- [22] Kader AA, Rolle RS (2004). In the role of post harvest management in assuring the quality and safety of horticultural produce, FAO Food and Agricultural Organizations of the United Nations, Rome.
- [23] Kutama AS, Aliyu BS, Muhammed I (2007). Fungal Pathogens Associated with Tomato Wicker Storage Baskets. *Science World Journal*, 2: 345-378
- [24] MC Kellar M.E. and E.B. Nelson 2003, compost induced suppression of Pythium damping off immediate by fatty acid metabolizing seed-colonizing microbial communities, *Applied and Environmental Microbiology* 69:452-460
- [25] MOI (2008). Ministry of Information, Sokoto, Nigeria. Diary. Pp. 12
- [26] Muhammad S, Shehu K, Amusa NA (2004). Survey of the market Diseases and aflatoxin contamination of tomato (*Lycopersicon esculentum* Mill.) Fruits in Sokoto, Northwestern Nigeria. *Nutrition and food science*, 34(276 Nigeria. Special FGN Gazette no. 23 on the 2006 Population Census. Patel NA, Dange SRS, Patel SI (2005). Efficacy of chemicals in controlling fruit rot of tomato caused by *Alternaria* tomato. *Indian J. Agric. Res.*, 39: 72-75.
- [27] Raj H and Kapoor 1997 gave possible management of Fusarium Wilt of tomato by soil amendment with compost.
- [28] Salau IA (2012). Studies of fungi associated with Human skin and vegetables disease in fadama land of Sokoto metropolis, Sokoto state, Nigeria. *M.Sc Thesis*, Usmanu Danfodiyo University, Sokoto, Nigeria
- [29] Salau IA, Kasarawa AB, Tambuwal NI (2012). Fungi associated with vegetable plant diseases in the Fadama land areas of Sokoto Metropolis. *Proceeding of Agricultural Society of Nigeria (ASN) 46th Annual Conference ("Kano 2012")* Pp: 779-782
- [30] Shehu K, Aliero AA (2010). Effects of purple blotch infection on the Proximate and mineral contents of onion Leaf *international Journal of Pharma Sciences and Research (IJPSR)* Vol.1(2), 131-133
- [31] Shehu K, Asma'u Muhammad M, Ibrahim AS (2014^b). A Preliminary study on Microbial Contamination of Leafy Vegetables in Sokoto Metropolis, Nigeria *Aceh Inter. J. Sci. and Technol.*, 3(3): 140-144
- [32] Shehu K, Muhammad S, Salau IA (2014^c). Fungi associated with fruit rot of tomato (*Solanum esculentum* Mill) in Sokoto. *American Journal of Research Communication*, 2014 www.usa-journals.com, ISSN: 2325-4076
- [33] Shehu K, Salau IA, Kasarawa AB, Umar AI (2014^a). Study on fungi associated with metropolitan waste water used for irrigation on soil in Sokoto metropolis. *Glo. Adv. Res. J. Agric. Sci.*, Vol. 3(6) pp. 148-151, June, 2014. Available online <http://garj.org/garjas/index.htm>
- [34] Tsoho BA (2004). Economics of Tomato Production under Small-scale Irrigation in Sokoto State. Unpublished M.Sc Thesis submitted to the Dept. of Agric. Economics and Farm Management. University of Ilorin, Ilorin, Nigeria.
- [35] Udo RK, Mamman AB (1993). *Nigeria: Giant in the tropics*. State surveys, pp. 435-446.



- [36] Veeken A.H., M.W.J. Blok, F.Curli, G.G.M. Coenen, A.J. Temorsheizer and H.V.M. Hamoleri 2005.
Improving quality of composted biowasted to enhance disease suppressiveness of compost amended
peat based potting mixes. *Soil biology and biochemistry* 37:2131-2140
- [37] Yar'adua AA (2003). Some studies on epidemiology of purple blotch of onion (*Allium cepa*) in Zaria,
Nigeria. *Danmasani* 1(6-8):98-106.