

Synthesis and spectroscopic studies of Ni(II) complexes of the tetraaza cyclohexadeca macrocyclic ligand.

* Kumari Anandi, **Dr. Ravindra Singh

*Research Scholar Dept. of Chemistry, Jai Prakash University Chapra, Saran Bihar

**Associate Prof. P.G. Dept. of Chemistry, Jai Prakash University Chapra, Saran Bihar

Abstract

In our present investigation, Synthesis and spectroscopic studies of Ni(II) complexes of the tetraaza cyclohexadeca macrocyclic ligand. The electronic spectra of Ni(II) complexes display a broad band in the charge transfer region indicative of their square planar geometry. The square planar geometry of Ni(II) complexes is further supported by their diamagnetic nature. Comparison of the infrared spectra of ligands and the complexes reveals facts. The appearance of a new band in IR spectra of $[NiL_1]Cl_2$, $[NiL_2]Cl_2$ and $[NiL_3]Cl_2$ at 610 cm^{-1} may be assigned to $\square Ni(II)$ mode of vibration. This also supports the coordination of the ligands to Ni(II) through nitrogen atoms. During the preparation of the complex of Ni(II) with 1,5,9,13-tetraaza-2,10-dimethyl-4,12-diphenyl cyclohexadeca-1,4,9,12-tetraene (L_1), Preparation of the complex of Ni(II) with 1,5,9,13-tetraaza-2,4,10,12-tetramethyl cyclohexadeca-1,4,9,12-tetraene (L_2) and Preparation of the complex of Ni(II) with 1,5,9,13-tetraaza-2,4,10,12-tetraphenyl cyclohexadeca-1,4,9,12-tetraene (L_3).

Keyword:- electronic spectra Ni(II), tetraaza, Dimethyl, diphenyl, tetraphenyltetramethyl, cyclohexadeca, tetraene.

INTRODUCTION

The chemistry of macrocyclic ligands has been a fascinating area of current research interest to the chemists all over the world. The continued interest and quest in designing new macrocyclic ligands stem mainly from their use as models for protein-metal binding sites in biological systems, as models for metalloenzymes, as sequestering reagents for specific metal ions, as models to study the magnetic exchange phenomena, as chemical sensors and batteries, as therapeutic reagents for the treatment of metal intoxication, as medical imaging agents, as catalysts, and in biomedical and fuel cell applications. The macrocycles


and their complexes have attracted the interest of both inorganic and bioinorganic chemists in recent years and have been the domain of their research work in view of their wide range of applications. The studies of macrocycles have undergone tremendous growth in past few decades and their complexation chemistry with a wide variety of metal ions has been extensively studied. Macrocyclic compounds are interesting ligand systems as they are good hosts for metal ions, neutral molecules and organic cation guests. The host-guest chemistry of metal- ions and macrocyclic compounds is quite significant in fundamental studies like phase-transfer catalysis. Macrocyclic complexes are considered to mimic the synthetic models of metalloporphyrins and metallocorrins due to their interinsic structural properties. Efforts have been made to achieve peripheral substitution, the appended substitution might create the possibility of synthesis of more complex compounds serving as new biologically important models. These compounds have received considerable attention due to their possible applications in medicine.

EXPERIMENTAL

PREPARATION OF LIGANDS

(i) Preparation of 1, 5, 9, 13-tetraaza-2,10-dimethyl- 4,12-diphenyl cyclohexadeca-1,4,9,12-tetraene (L₁):

0.10 mole of 1,3-diaminopropane was dissolved in 50 ml. of methanol and cooled in ice. To this solution, a solution of 0.10 mole of 1-phenylbutane-1,3-dione in 50 ml. of methanol was added. The mixture was stirred for 24 hours at 5°C, refluxed for about 8 hours and then concentrated to about 30 ml. and placed in an ice-bath. White coloured crystals were separated which were washed with ice-cold methanol and recrystallised twice from methanol- water mixture.





(Fig. -1)

(ii) Preparation of 1,5,9,13-tetraaza-2,4,10,12- tetramethyl cyclohexadeca-1,4,9,12-tetraene (L₂)

0.10 mole of 1,3-diaminopropane was dissolved in 50 ml. of methanol and cooled in an ice-bath. A solution of 0.10 mole of pentane-2,4-dione in 50 ml. of methanol was added to the methanolic solution of 1,3-diaminopropane. The mixture was stirred for about 24 hours at 5°C, refluxed for about 8 hours and then concentrated to about 30 ml. and placed in an ice-bath. Yellowish-white crystals were separated which were washed with ice-methanol

and recrystallised twice from methanol-watermixture.




1,5,9,13-tetraaza-2,4,10,12-tetramethyl cyclohexadeca- 1,4,9,12-tetraene(L₂)

(Fig. -2)


(iii) Preparation of 1,5,9,13-tetraaza-2,4,10,12- tetraphenyl cyclohexadeca-1,4,9,12-tetraene (L₃)

0.10 mole of 1,3-diamino propane was dissolved in 50 ml. of methanol and cooled in an ice-bath. A solution of 0.10 mole 1,3-diphenylpropane-1,3-dione in 50 ml..of methanol was

prepared and added to the methanolic solution of 1,3-diaminopropane. The mixture was stirred for about 24 hours at 5°C, refluxed for about 8 hours and then concentrated to about 30 ml. and placed in an ice-bath washed with ice-cold methanol and re-crystallized twice from methanol-watermixture.



1,3-diphenyl propane-1,3-dione



1,5,9,13-tetraaza-2,4,10,12-tetraphenyl cyclohexadeca-1,5,9,12-tetraene(L₃)

Fig. -3

MICROANALYTICAL DATA RELATED TO LIGANDS :

Table-1

Sl. No.	Ligands	% Found (% Calculated)			Molecular Formula	Colour
		C	H	N		
1.	L ₁	79.28 (78.00)	7.92 (8.00)	13.84(14.00)	C ₂₆ H ₃₂ N ₄	White
2.	L ₂	69.82 (69.56)	10.26 (10.14)	20.29 (20.28)	C ₁₆ H ₂₈ N ₄	Yellowish White
3.	L ₃	82.21 (82.44)	6.74 (6.87)	10.53 (10.88)	C ₃₆ H ₃₆ N ₄	White

Microanalytical data of ligands are in confirmity with the molecular formula of corresponding ligands.

PREPARATION OF COORDINATION COMPOUNDS

As the yield of the macrocyclic ligands were very low, their complexes with Ni (II), were prepared by template method.

(i) Preparation of the complex of Ni (II) with 1,5,9,13- tetraaza-2,10-dimethyl-4,12-diphenylcyclohexadeca- 1,4,9,12-tetraene (L₁):

0.10 mole of NiCl₂.6H₂O was dissolved in 50 ml. of water-methanol mixture (1:1 volume ratio). It was taken in a round-bottom flask of 500 ml. capacity. To this solution were added 50 ml. methanolic solution of 0.20 mole of 1,3-diaminopropane and 50 ml. methanolic solution of 0.20 mole of 1-phenylbutane-1,3-dione. The mixture was refluxed on a sand-bath using water condenser for about 8 hours and then left for about 24 hours. Green crystals were separated out which were filtered, washed with water and then with methanol. The product was dried in an electric oven at 110°C. The product was insoluble in water, methanol, ethanol, CHCl₃ and CCl₄ but it was soluble in DMF and DMSO. The m.p. of the complex was recorded to be 212°C.

(ii) Preparation of the complex of Ni (II) with 1,5,9,13- tetraaza-2,4,10,12-tetramethyl cyclohexadeca-1,4,9,12- tetraene (L₂):

0.10 mole of NiCl₂.6H₂O was dissolved in 50 ml. of water-methanol mixture (1 : 1 volume ratio). It was taken in a round-bottom flask of 500 ml. capacity. To this solution were added 50 ml. methanolic solution of 0.20 mole of 1,3-diaminopropane and 50 ml. methanolic solution of 0.20 mole of pentane-2,4-dione. The mixture was refluxed on a water-bath using water-condenser for about 8 hours and then left for about 24 hours. Green crystals were separated out which were filtered, washed with water and then with methanol. The product was dried in an electric oven at 110°C. It was insoluble in water and common organic solvents such as methanol, ethanol, benzene, ether, CHCl₃ and CCl₄ but was soluble in DMF and DMSO. The m.p. of the complex was recorded to be 204°C.

(iii) Preparation of the complex of Ni (II) with 1,5,9,13- tetraaza-2,4,10,12-tetraphenyl cyclohexadeca-1,4,9,12- tetraene (L₃):

0.10 mole of NiCl₂.6H₂O was dissolved in 50 ml. of water-methanol mixture (1:1 volume ratio). It was taken in a round-bottom flask of 500 ml. capacity. To this solution were added 50 ml. of methanolic solution of 0.20 mole of 1,3-diaminopropane and 50 ml. methanolic solution of 0.20 mole of 1,3-diphenylpropane-1,3-dione. The mixture was refluxed on a water-bath using a water condenser for about 8 hours. It was then allowed to cool for another 24 hours. Green crystals were separated which were filtered, washed with water and then with methanol. The complex was insoluble in water and common organic solvents such as benzene, ether, methanol, ethanol, CHCl₃, CCl₄. It was soluble in DMF and DMSO. The complex was dried in an electric furnace at 110°C and its m.p. was recorded to be 119°C.

Result & Discussions

Spectroscopic Investigations of Complexes

Electronic spectra of complexes of Ni (II):

The free ion ground term of Ni (II) is 3F with 3P as the first excited term. Under the influence of the cubic field, the 3F term splits into ${}^3T_{1g}(F)$, ${}^3T_{2g}(F)$ and ${}^3A_{2g}(F)$ states and the 3P term transforms to ${}^3T_{1g}(P)$ state only. Ni(II) complexes in weak octahedral field, are expected to give three d-d bands corresponding to the transitions, ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F)$, ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F)$ and ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(P)$. In case of Ni(II) tetrahedral complexes, three d-d transition bands are also expected corresponding to the transitions ${}^3T_1 \rightarrow {}^3T_2$, ${}^3T_1 \rightarrow {}^3E_2$ and ${}^3T_1 \rightarrow {}^3T_1(P)$. Distorted octahedral complexes of Ni (II) are expected to give four d-d transition bands corresponding to the transitions ${}^3B_{1g} \rightarrow {}^3E_g$, ${}^3B_{1g} \rightarrow {}^3B_{2g}$, ${}^3B_{1g} \rightarrow {}^3T_{1g}({}^3E_g, {}^3A_{2g})$ and ${}^3B_{1g} \rightarrow {}^3T_{1g}(P)$. Low spin Ni(II) square planar complexes should give two d-d transition bands but there occur at high energy in the charge transfer region and have not been satisfactorily interpreted. In high spin square planar complexes of Ni(II), two transitions corresponding to ${}^1A_{1g} \rightarrow {}^1B_{2g}$ and ${}^1A_{1g} \rightarrow {}^1E_g$ are expected. In our present investigation, the electronic spectra of Ni(II) complexes display a broad band in the charge transfer region indicative of their square planar geometry. The square planar geometry of Ni(II) complexes is further supported by their diamagnetic nature.

Infrared Spectra of complexes:

Comparison of the infrared spectra of ligands and the complexes reveals following facts :

- (i) The appearance of a new band in IR spectra of $[NiL_1]Cl_2$, $[NiL_2]Cl_2$ and $[NiL_3]Cl_2$ at 610 cm^{-1} may be assigned to ν_{Ni-N} mode of vibration¹⁹³. This also supports the coordination of the ligands to Ni(II) through nitrogen atoms.

^1H NMR spectra of complexes:

^1H NMR spectra of complexes further substantiate the proposed structure of complexes.

The ^1H NMR spectral data of complexes are given in Table -2.

Table- 2

^1H NMR Spectral data of complexes

Sl.No.	Complex	Position of ^1H NMR signals Compound
1.	$[\text{NiL}_1]\text{Cl}_2$	$\square = 0.90$ ppm (s, 6H), $\square = 1.2$ ppm (m, 12H), $\square = 1.7$ ppm (s, 4H), $\square = 7.5$ ppm (m, 10H)
2.	$[\text{NiL}_2]\text{Cl}_2$	$\square = 1.0$ ppm (s, 12H), $\square = 1.4$ ppm (m, 12H), $\square = 1.8$ ppm (s, 4H)
3.	$[\text{NiL}_3]\text{Cl}_2$	$\square = 1.2$ ppm (m, 12H), $\square = 1.5$ ppm (s, 4H), $\square = 7.2$ ppm (m, 20H)

s = singlet, m = multiplet.

The ^1H NMR signals at $\square = 0.90$ ppm are due to methyl ($-\text{CH}_3$) protons. The multiplet bands at $\square = 1.2$ ppm to 1.4 ppm are due to methylene ($-\text{CH}_2$) protons of $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ moiety. The singlet signals at $\square = 1.5$ ppm to 1.8 ppm are due to the allylic protons of methylene ($-\text{CH}_2-$) groups. The multiplet signals centered at $\square = 7.2$ ppm to 7.5 ppm are due to the aromatic protons of phenyl groups.

CONCLUSION

Macrocyclic compounds find useful applications:

- (i) In removing heavy metals from aqueous solution for water purification.
- (ii) As molecular switches and linear motors for constructing artificial nanoscale machinery (rotaxanes).
- (iii) As chemical sensors.
- (iv) In mimicry of cellular receptors.

(v) As molecular recognition agents.

(vi) As recognition agent for peptides.

(vii) As organic light emitting diodes.

Keeping in view the wide range of applications of macrocyclic compounds and their complexes, we undertook a programme to synthesise and characterize complexes of Ni(II), Pd(II) and Pt(II) with 1,5,9,13-tetraaza-2,10-dimethyl-4,12-diphenylcyclohexadeca-1,4,9,12-tetraene (L_1), 1,5,9,13-tetraaza-2,4,10,12-tetramethylcyclohexadeca-1,4,9,12-tetraene (L_2) and 1,5,9,13-tetraaza-2,4,10,12-tetraphenylcyclohexadeca-1,4,9,12-tetraene (L_3).

REFERENCES

1. Ashu Chaudhary Singh:- *Rasayan J. Chem.*, 2 (1), 191 and R. V.S (2009).
2. B.D.Busch:- *Rec. Chem. Progr.*; **25**, 107(1964).
3. Bartlett et al.:- *J. Am. Chem. Soc.*; **117**10025 (1995).
4. C.N.Schrauzer:- *Chem. Ber*; **95**, 1438(1962).
5. D. R. Boston and N. j. Rose:- *Abstracts, 157th National Meetings of the American Chemical Society, Minneapolis, Minn, April 1969, No. INOR-96.*
6. L. T. Bozic, E. Marotta, P. Taraldi:- *Polyhedron*; **26**, 1663 (2007).
7. Levason et al.:- *Inorg. Chem.*; **30**, 331(1991); **34**, 1626(1995)
8. M. B. Reddy, K. Shanker, P. Usha Rani, R. Rohini, K. Reddy, V. Ravinder. :- *J. Indian Chem. Soc.*, **4**, 971 (2007) Ch.
9. Mayur A. Panchbhai, L. J. Paliwal and N. S. Bhave:- *E-Journal of chemistry*; **5**(S1), 1048(2008).
10. P. M. Reddy, A. V. V. S. Prasad, R. Rohini, V. Ravinder:- *Spectrochim. Acta.*, **70A**, 704(2008)
11. P. M. Reddy, Y. P. HO, K. Shankar, R. Rohini, V. Ravinder. :- *Eur. J. Med. Chem.*; **10**, 1(2009)
12. S. Chandra, A. Gautam, M. Tyagi:- *Transition Met. Chem.*, **32**, 1079(2007)
13. S. Ilhan, H. Temel.:- *Transition Met. Chem.*, **32**, 1039(2007)